HyQvia® (Immune Globulin SubQ Infusion) Referral Form





PATIENT INFORMATION	Referral Status:	New Referral	Updated Referral	Referral Renewal
DOB: Patient Name:			Patient Phone:	
Patient Address:			Patient Email:	
NKDA Allergies:			Weight (lbs/kg):	Height:
ICD-10 Code (required): ICD-10 Description:	Last Treatme	ent Date:	Last 4 Dig	gits SSN:
PROVIDER INFORMATION				
Referral Coordinator Name:	Referral Coo	rdinator Email:		
Ordering Provider:	Provider NPI	:		
Referring Practice Name:	Phone:		Fax:	
Practice Address:	City:		State: Z	Zip Code:
Physician Preferred Method of Contact: Email:	F	ax:	Pho	ne:
NURSING ☐ Infusion to be administered per Vivo protocols. LABORATORY ORDERS CBC at each dose every	Patronidase to I Pharmacist will ca Pat tree intr Pat Sub Wee	For ient switching fratment: Administe avenous treatment ient naïve to IgG trocutaneous (Humanek intervals, after in Dose: Interval: witching from IVIG	PI Patients Only rom Immune Globul r Hyqvia at the same d t, after initial ramp-up. reatment or switching n) [IGSC]: Administer H nitial ramp up.* For CIDP Patients Onl (human) treatment, ad	*Ramp-up schedule on page 2. lin Intravenous (Human) [IVIII] lose and frequency as the previor * from Immune Globulin lyqvia at 300 to 600 mg/kg at 3 to
REQUIRED DOCUMENTATION Patient Demographics Insurance Card/Information Progress Notes Supporting DX Medication List and H&P Serum Creatinine (within last 3 months if treatment naive) **Consider administering premedication for prophylaxis against infusion in	Additional I	Interval: Notes	**Order is valid for one	e year unless otherwise noted**
Provider Name (Print) Provider	r Signature			Date
Email Referrals To: referrals@vivoinfusion.com OR Fax Below	, Ha	ve a Question?	Call (720) 902-411:	1

 Email Referrals To: referrals@vivoinfusion.com OR Fax Below
 Have a Question? Call (720) 902-4111

 Colorado: 303-418-4679
 Michigan: 833-957-2188
 New York: 800-540-1852
 Texas: 469-340-0044

 Connecticut: 203-724-4838
 Minnesota: 763-290-0903
 Ohio: 216-400-0674
 Virginia: 804-500-5941

 Florida: 904-930-4211
 Nevada: 702-489-5744
 Oklahoma: 918-770-4421
 Wisconsin: 414-600-5383

 Massachusetts: 781-202-1629
 New Jersey: 609-955-3711
 Pennsylvania: 215-399-9244

Ramp up Schedule for Hyqvia

Initial Treatment Interval and Ramp-Up Schedule for PI

For patients previously on another IgG treatment, the first dose should be given approximately 1 week after the last infusion of their previous treatment.

PI: Ramp-up schedule if switching from IVIG

Week	Dose Interval	Dose
1	1st Dose	Total grams x 0.25
2	2nd Dose	Total grams x 0.50
3	NO INFUSION	NO INFUSION
4	3rd Dose	Total grams x 0.75
5	NO INFUSION	NO INFUSION
6	NO INFUSION	NO INFUSION
7	4th Dose	Total grams

PI: Ramp-up schedule if switching from SCIG

Treatment Interval	Dosing Q 4 Weeks	Dosing Q3 Weeks
1st Infusion (Week 1)	Grams x 0.25	Grams x 0.33
2nd Infusion (Week 2)	Grams x 0.5	Grams x 0.67
3rd Infusion (Week 4)	Grams x 0.75	Administer Total Grams
4th Infusion (Week 7)	Administer Total Grams	N/A

Initial Treatment Interval and Ramp-Up Schedule for CIDP

Doses less than or equal to 0.4 g/kg can be administered without ramp-up Patients must be on stable doses of IVIG for 12 weeks before switching to Hyqvia

Week	Dose Interval	Dose
	Switch from IVIG	
1	No Infusion	
2	1st Dose	Total grams x 0.25
3	2nd Dose	Total grams x 0.25
4	3rd Dose	Total grams x 0.50
6	4th Dose	Total grams x 0.75
9	5th Dose	Total grams

Total grams=total monthly equivalent dose in grams.