

# HyQvia® (Immune Globulin SubQ Infusion) Referral Form



**Preferred Clinic** (select one):

## PATIENT INFORMATION

New Referral

Updated Referral

Referral Renewal

DOB:	Patient Name:	Patient Phone:
Patient Address:	Patient Email:	
NKDA Allergies:	Weight (lbs/kg):	Height:
ICD-10 Code (required):	ICD-10 Description:	Last Treatment Date:
		Last 4 Digits SSN:

## PROVIDER INFORMATION

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:
Physician Preferred Method of Contact:	Email:	Fax:	Phone:

## STANDING ORDERS

☒ Infusion to be administered per Vivo protocols.

## LABORATORY ORDERS

CBC every \_\_\_\_\_  
CMP every \_\_\_\_\_  
OTHER \_\_\_\_\_

## PREMEDICATIONS

acetaminophen (Tylenol) 500 mg 650 mg 1000 mg PO  
cetirizine (Zyrtec) 10mg PO  
loratadine (Claritin) 10mg PO  
diphenhydramine (Benadryl) 25 mg 50 mg PO IV  
methylprednisolone (Solu-Medrol) 40mg 125mg IV  
hydrocortisone (Solu-Cortef) 100mg IV  
Other: \_\_\_\_\_  
Dose: \_\_\_\_\_ Route: \_\_\_\_\_

## REQUIRED DOCUMENTATION

Patient Demographics

Insurance Card/Information

Progress Notes Supporting DX

Medication List and H&P

Serum Creatinine (within last 3 months if treatment naive)

**\*\*Consider administering premedication for prophylaxis against infusion reactions and hypersensitivity reactions. \*\*Order is valid for one year unless otherwise noted\*\***

## HYQVIA ADMINISTRATION

*Hyaluronidase to infuse first at 1-2 ml/minute/site subcutaneous administration.*

*Pharmacist will calculate ramp-up schedule unless otherwise noted. \*Ramp-up schedule on page 2.*

### For PI Patients Only

**Patient switching from Immune Globulin Intravenous (Human) [IVIG] treatment:** Administer Hyqvia at the same dose and frequency as the previous intravenous treatment, after initial ramp-up.\*

**Patient naïve to IgG treatment or switching from Immune Globulin Subcutaneous (Human) [IGSC]:** Administer Hyqvia at 300 to 600 mg/kg at 3 to 4 week intervals, after initial ramp up.\*

Dose: \_\_\_\_\_  
Interval: \_\_\_\_\_

### For CIDP Patients Only

If switching from IVIG (human) treatment, administer Hyqvia at the same dose and frequency as the previous IV treatment, after the initial dose ramp-up.\*

Dose: \_\_\_\_\_  
Interval: \_\_\_\_\_

## Additional Notes

Provider Name (Print)	Provider Signature	Date
-----------------------	--------------------	------

Email Referrals To: [referrals@vivoinfusion.com](mailto:referrals@vivoinfusion.com) OR Fax Below

Have a Question? Call (720) 902-4111

Colorado: 303-418-4679	Michigan: 833-957-2188	New York: 800-540-1852	Texas: 469-340-0044
Connecticut: 203-724-4838	Minnesota: 763-290-0903	Ohio: 216-400-0674	Virginia: 804-500-5941
Florida: 904-930-4211	Nevada: 702-489-5744	Oklahoma: 918-770-4421	Wisconsin: 414-600-5383
Massachusetts: 781-202-1629	New Jersey: 609-955-3711	Pennsylvania: 215-399-9244	

# Ramp up Schedule for Hyqvia

## Initial Treatment Interval and Ramp-Up Schedule for PI

For patients previously on another IgG treatment, the first dose should be given approximately 1 week after the last infusion of their previous treatment.

### PI: Ramp-up schedule if switching from IVIG

Week	Dose Interval	Dose
1	1st Dose	Total grams x 0.25
2	2nd Dose	Total grams x 0.50
3	NO INFUSION	NO INFUSION
4	3rd Dose	Total grams x 0.75
5	NO INFUSION	NO INFUSION
6	NO INFUSION	NO INFUSION
7	4th Dose	Total grams

### PI: Ramp-up schedule if switching from SCIG

Treatment Interval	Dosing Q 4 Weeks	Dosing Q3 Weeks
1st Infusion (Week 1)	Grams x 0.25	Grams x 0.33
2nd Infusion (Week 2)	Grams x 0.5	Grams x 0.67
3rd Infusion (Week 4)	Grams x 0.75	Administer Total Grams
4th Infusion (Week 7)	Administer Total Grams	N/A

## Initial Treatment Interval and Ramp-Up Schedule for CIDP

Doses less than or equal to 0.4 g/kg can be administered without ramp-up

Patients must be on stable doses of IVIG for 12 weeks before switching to Hyqvia

Week	Dose Interval	Dose
	Switch from IVIG	
1	No Infusion	
2	1st Dose	Total grams x 0.25
3	2nd Dose	Total grams x 0.25
4	3rd Dose	Total grams x 0.50
6	4th Dose	Total grams x 0.75
9	5th Dose	Total grams

Total grams=total monthly equivalent dose in grams.